

Bibliografia consultata

- Arapitsas, P., et al. (2021). Improving the phloroglucinolysis protocol and characterization of sagrantino wines proanthocyanidins. *Molecules*, 26(4), 1087. <https://doi.org/10.3390/molecules26041087>
- Bindon, K., et al. (2014). Relationships between harvest time and wine composition in *Vitis vinifera* cv. Cabernet Sauvignon 2. *Food Chemistry*, 154, 90-101. <https://doi.org/10.1016/j.foodchem.2013.12.099>
- Brossaud, F., Cheynier, V., & Noble, A. C. (2001). Bitterness and astringency of grape and wine polyphenols. *Australian Journal of Grape and Wine Research*, 7(1), 33-39. <https://doi.org/10.1111/j.1755-0238.2001.tb00191.x>
- Castellarin, S. D., et al. (2007). Water deficits accelerate ripening and induce changes in gene expression regulating flavonoid biosynthesis in grape berries. *Planta*, 227(1), 101-112. <https://doi.org/10.1007/s00425-007-0598-8>
- Cejudo-Bastante, M. J., et al. (2011). Micro-oxygenation and oak chip treatments of red wines. *Food Chemistry*, 124(3), 727-737. <https://doi.org/10.1016/j.foodchem.2010.06.087>
- Cheynier, V., et al. (2006). Structure and properties of wine pigments and tannins. *American Journal of Enology and Viticulture*, 57(3), 298-305. <https://doi.org/10.5344/ajev.2006.57.3.298>
- Cortell, J. M., et al. (2007). Influence of vine vigor on grape anthocyanins. *Journal of Agricultural and Food Chemistry*, 55(16), 6575-6584. <https://doi.org/10.1021/jf070195v>
- Del Barrio-Galan, R., et al. (2015). Relationship between metal content in bottled aged red wines and changes in phenolic composition. *Food Research International*, 77, 304-313. <https://doi.org/10.1016/j.foodres.2015.08.028>
- Delgado, R., et al. (2004). Changes in the phenolic composition of grape berries during ripening in relation to vineyard nitrogen and potassium fertilisation rates. *Journal of the Science of Food and Agriculture*, 84(7), 623-630. <https://doi.org/10.1002/jsfa.1685>
- Evyugin, D. D., Magina, S., & Evtuguin, D. V. (2020). Recent advances in the production and applications of ellagic acid and its derivatives. A review. *Molecules*, 25(12), 2745. <https://doi.org/10.3390/molecules25122745>

- Fernandez de Simon, B., et al. (2010). Effect of size, seasoning and toasting in the volatile compounds in toasted oak wood. *Analytica Chimica Acta*, 660(1-2), 211-220. <https://doi.org/10.1016/j.aca.2009.09.031>
- Fulcrand, H., et al. (2006). Phenolic reactions during winemaking and aging. *American Journal of Enology and Viticulture*, 57(3), 289-297. <https://doi.org/10.5344/ajev.2006.57.3.289>
- Gutierrez-Escobar, R., Aliano-Gonzalez, M. J., & Cantos-Villar, E. (2021). Wine Polyphenol Content and Its Influence on Wine Quality and Properties: A Review. *Molecules*, 26(3), 718. <https://doi.org/10.3390/molecules26030718>
- Hanlin, R. L., et al. (2010). Condensed tannin and grape cell wall interactions and their impact on tannin extractability into wine. *Australian Journal of Grape and Wine Research*, 16(1), 173-188. <https://doi.org/10.1111/j.1755-0238.2009.00068.x>
- He, F., et al. (2012). Anthocyanins and their variation in red wines I. Monomeric anthocyanins and their color expression. *Molecules*, 17(2), 1571-1601. <https://doi.org/10.3390/molecules17021571>
- Kennedy, J. A., & Jones, G. P. (2001). Analysis of proanthocyanidin cleavage products following acid-catalysis in the presence of excess phloroglucinol. *Journal of Agricultural and Food Chemistry*, 49(4), 1740-1746. <https://doi.org/10.1021/jf0010300>
- Kennedy, J. A., Matthews, M. A., & Waterhouse, A. L. (2002). Effect of maturity and vine water status on grape skin and wine flavonoids. *American Journal of Enology and Viticulture*, 53(4), 268-274.
- Martinez-Gil, A., Cadahía, E., Fernández de Simón, B., & Nevares, I. (2019). Phenolic and volatile compounds in *Quercus humboldtii* Bonpl. wood: effect of toasting with respect to oaks traditionally used in cooperage. *Journal of the Science of Food and Agriculture*, 99(1), 315-324. <https://doi.org/10.1002/jsfa.9190>
- Monagas, M., et al. (2005). Updated knowledge about the presence of phenolic compounds in wine. *Critical Reviews in Food Science and Nutrition*, 45(2), 85-118. <https://doi.org/10.1080/10408690490911710>
- Nevares, I., et al. (2021). Characterization of the oxygen transmission rate of new-ancient natural materials for wine maturation containers. *Foods*, 10(1), 140. <https://doi.org/10.3390/foods10010140>

- Nevarés, I., et al. (2021). Characterization of the oxygen transmission rate of new-ancient natural materials for wine maturation containers. *Foods*, 10(1), 140. <https://doi.org/10.3390/foods10010140>
- Ribereau-Gayon, P., et al. (2006). *Handbook of Enology, Vol. 2: The Chemistry of Wine Stabilization and Treatments* (2nd ed.). John Wiley & Sons.
- Ristic, R., & Iland, P. G. (2005). Relationships between seed and berry development of *Vitis vinifera* cv Shiraz. *Australian Journal of Grape and Wine Research*, 11(1), 43-58. <https://doi.org/10.1111/j.1755-0238.2005.tb00278.x>
- Rodriguez-Bencomo, J. J., et al. (2009). Volatile compounds of red wines macerated with Spanish, American, and French oak chips. *Journal of Agricultural and Food Chemistry*, 57(14), 6383-6391. <https://doi.org/10.1021/jf900739k>
- Sacchi, K. L., Bisson, L. F., & Adams, D. O. (2005). A review of the effect of winemaking techniques on phenolic extraction in red wines. *American Journal of Enology and Viticulture*, 56(3), 197-206.
- Van Leeuwen, C., et al. (2009). Vine water status is a key factor in grape ripening and vintage quality. *Journal International des Sciences de la Vigne et du Vin*, 43(3), 121-134. <https://doi.org/10.20870/oeno-one.2009.43.3.798>
- Vivas, N., & Glories, Y. (1996). Role of oak wood ellagitannins in the oxidation process of red wines during aging. *American Journal of Enology and Viticulture*, 47(1), 103-107.
- WatreLOT, A. A., & Norton, E. L. (2020). Chemistry and reactivity of tannins in *Vitis* spp.: a review. *Molecules*, 25(9), 2110. <https://doi.org/10.3390/molecules25092110>
- Yilmaz, Y., & Toledo, R. T. (2004). Major flavonoids in grape seeds and skins: antioxidant capacity of catechin, epicatechin, and gallic acid. *Journal of Agricultural and Food Chemistry*, 52(2), 255-260. <https://doi.org/10.1021/jf030117h>